

Supersymmetric Model with Grassmann-odd Lagrangian

D.V. Soroka^a, V.A. Soroka^b and J. Wess^{c, d}

^aApt. 48, Val'ter Street 7, 61108 Kharkov, Ukraine

^bInstitute for Theoretical Physics, NSC "Kharkov Institute of Physics and Technology",
61108 Kharkov, Ukraine

^cSektion Physik, Universität München, Theresienstr. 37, D-80333 München

^dMax-Planck-Institut für Physik, Föhringer Ring 6, D-80805 München

A supersymmetric $D = 1, N = 1$ model with a Grassmann-odd Lagrangian is proposed which, in contrast to the model with an even Lagrangian, contains not only a kinetic term but also an interaction term for the coordinates entering into one real scalar Grassmann-even (bosonic) superfield.

1. Introduction

There are two ways for the formulation of Hamiltonian dynamics in phase superspace: either by means of an even Poisson bracket with the help of a Grassmann-even Hamiltonian or in an odd Poisson bracket [1] with a Grassmann-odd Hamiltonian¹. The dynamical systems with equal numbers of pairs of even and odd phase coordinates admit equivalent descriptions in the both brackets at once [2,3] with the help of the corresponding equivalent Hamiltonians of opposite Grassmann parities. However, there are such Hamiltonian systems which permit for their description only the bracket of a definite parity. For example, the Hamilton systems with odd equal numbers of the even and odd canonical variables can be described only by means of the odd Poisson bracket (see, for example, [4]). On the other hand, the systems having an even number of the even canonical variables which differs from a number of the odd variables allow their Hamiltonian description only in terms of the even Poisson bracket. In this connection, by assuming that a Hamilton system has the corresponding to it Lagrangian formulation and taking into account that in this case the Grassmann parity of the Lagrangian coincides with the parities of the bracket

and Hamiltonian in terms of which this Hamiltonian system is described, we see that there are also two ways for the formulation of Lagrangian dynamics in configuration superspace: either by means of an even Lagrangian or in terms of an odd one [3,5,6]. As in the case of the Hamiltonian dynamics, there are such systems which can be described only with the help of the Grassmann-odd Lagrangian and have no description in terms of the even one and vice versa.

In this report we just give such examples for $D = 1, N = 1$ supersymmetric models. By using the formalism with the even Lagrangian, we can construct from a real bosonic scalar superfield only kinetic terms, but a term for interaction turns out to be a total time derivative. While, using the formalism with the odd Lagrangian, we are able to construct from this superfield a model which contains both the kinetic and interaction terms.

The report is organized as follows. In Section 2 we show that in the case $D = 1, N = 1$ supersymmetry the even Lagrangian, which reduced from Lagrangian density constructed from one real scalar bosonic superfield, does not contain an interaction term. Then we propose a $D = 1, N = 1$ supersymmetric model having the odd Lagrangian, which also constructed on the basis of this superfield but has an interac-

¹We shall only consider non-degenerate brackets.

tion term. Consideration in this Section is carried out in the Lagrangian component formulation. The Hamiltonian component formulation for this model is given in Section 3. In Section 4 we present a superfield approach for the model both in the Lagrangian and Hamiltonian formalisms. In Section 5 we show once again that even in a particular case of the supersymmetric one-dimensional oscillator the even Lagrangians, which are dynamically equivalent to the model proposed, can not be reduced from any superfield Lagrangian densities constructed from one real bosonic $D = 1, N = 1$ superfield.

2. Lagrangian component formulation

With respect to $D = 1, N = 1$ supersymmetry for the proper time t and its real Grassmann superpartner η

$$t' = t + i\epsilon\eta, \quad \eta' = \eta + \epsilon$$

the components of the real Grassmann-even (bosonic) $(g(\Phi) = 0)^2$ scalar superfield

$$\Phi(t, \eta) = q(t) + i\eta\theta(t) = \Phi'(t', \eta') \quad (1)$$

are transformed as

$$\begin{aligned} \delta q(t) &\equiv q'(t) - q(t) = i\theta\epsilon, \\ \delta\theta(t) &\equiv \theta'(t) - \theta(t) = \dot{q}\epsilon, \end{aligned} \quad (2)$$

where a dot means a time derivative and δ is a variation of the form.

With the help of a covariant derivative

$$D = \partial_\eta - i\eta\partial_t, \quad (3)$$

where $\partial_x \equiv \frac{\partial}{\partial x}$, we can construct from Φ an invariant action S_0 with a Grassmann-even Lagrangian l , $(g(l) = 0)$

$$S_0 = \int dt d\eta \left[\frac{i}{2} D\Phi\dot{\Phi} + iDV(\Phi) \right] = \int dt l^0,$$

which contains only kinetic terms for the bosonic q and fermionic θ coordinates

$$l^0 = \frac{\dot{q}^2}{2} + \frac{i}{2}\theta\dot{\theta} + \dot{V}(q),$$

² $g(\Phi)$ is a Grassmann parity of the quantity Φ .

while their interaction term with an arbitrary real function $V(\Phi)$ of Φ is a total time derivative.

By using a Grassmann-odd Lagrangian $\overset{1}{L}$ ($g(\overset{1}{L}) = 1$) which in the case of $D = 1, N = 1$ supersymmetry corresponds to an even Lagrangian density $\mathcal{L}$ ($g(\mathcal{L}) = 0$), we are able to construct a supersymmetric action

$$\begin{aligned} S_1 &= -i \int dt d\eta \left[\frac{\dot{\Phi}^2}{2} - V(\Phi) \right] = -i \int dt d\eta \mathcal{L} \\ &= \int dt \overset{1}{L}, \end{aligned} \quad (4)$$

including both a kinetic term for coordinates q, θ and a term describing their interaction with the help of an arbitrary real function $V(\Phi)$. The Lagrangian density superfield has the form

$$\mathcal{L} = \overset{0}{L} + i\eta\overset{1}{L} = \frac{\dot{q}^2}{2} - V(q) + i\eta[\dot{q}\dot{\theta} - \theta V'(q)], \quad (5)$$

where a prime denotes a derivative with respect to the coordinate q . Note that the even and odd Lagrangians $\overset{0}{L}$ and $\overset{1}{L}$ are connected with each other with the aid of the exterior differential³ d that accompanied by the following map $\lambda : dq \rightarrow \theta$ (see, for example, [6])

$$\lambda : d\overset{0}{L} \rightarrow \overset{1}{L}$$

and are transformed under supersymmetry (2) as

$$\delta\overset{0}{L} = i\overset{1}{L}\epsilon, \quad \delta\overset{1}{L} = \dot{\overset{0}{L}}\epsilon.$$

In accordance with the scheme given in [6], the Euler-Lagrange equation for the even Lagrangian $\overset{0}{L}$ coincides with that one for the odd Lagrangian $\overset{1}{L}$, which corresponds to the variable θ ,

$$\begin{aligned} \frac{d}{dt} \left(\partial_{\dot{\theta}} \overset{1}{L} \right) - \partial_{\theta} \overset{1}{L} &= \frac{d}{dt} \left(\partial_{\dot{q}} \overset{0}{L} \right) - \partial_q \overset{0}{L} \\ &= \ddot{q} + V'(q) = 0, \end{aligned} \quad (6)$$

while the Euler-Lagrange equation for $\overset{1}{L}$, corresponding to the variable q , can be obtained by

³We adopt the Grassmann parity of the exterior differential to be equal to unit $g(dx) = g(x) + 1$.

taking the exterior differential from two last equations in (6) and performing the map λ

$$\begin{aligned} \frac{d}{dt} \left(\partial_{\dot{q}} dL \right) - \partial_q dL &= d\ddot{q} + dqV'' = 0 \\ \xrightarrow{\lambda} \frac{d}{dt} \left(\partial_{\dot{q}} \overset{1}{L} \right) - \partial_q \overset{1}{L} &= \ddot{\theta} + \theta V'' = 0 . \end{aligned} \quad (7)$$

3. Hamiltonian component formulation

The Lagrangians $\overset{0}{L}$ and $\overset{1}{L}$ in (5) have the same even momentum

$$p = \partial_{\dot{q}} \overset{0}{L} = \partial_{\dot{\theta}} \overset{1}{L} = \dot{q} ,$$

whereas an odd momentum π for $\overset{1}{L}$ is connected with it by means of the exterior differential d and the map λ

$$\lambda : dp \rightarrow \pi = \partial_{\dot{q}} \overset{1}{L} = \dot{\theta} .$$

Corresponding to the even Lagrangian $\overset{0}{L}$ an even Hamiltonian

$$\overset{0}{H} = \dot{q}p - \overset{0}{L} = \frac{p^2}{2} + V(q) \quad (8)$$

is also related with an odd Hamiltonian $\overset{1}{H}$, responding to the odd Lagrangian $\overset{1}{L}$,

$$\overset{1}{H} = \dot{q}\pi + \dot{\theta}p - \overset{1}{L} = p\pi + \theta V'(q) \quad (9)$$

in a similar way

$$\lambda : d\overset{0}{H} \rightarrow \overset{1}{H} .$$

In the case of Hamiltonian formulation, the scheme given in Refs. [6,7] prescribe that the Hamilton equations

$$\dot{q} = p , \quad \dot{p} = -V'(q) \quad (10)$$

formulated for the even phase coordinates $x^i = (q, p)$ with the help of the even Hamiltonian (8) in the even Poisson bracket

$$\dot{x}^i = (x^i, \overset{0}{H})_0 \equiv x^i \left(\overleftarrow{\partial}_q \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_q \right) \overset{0}{H} \quad (11)$$

coincide with those ones obtained for x^i by using the odd Hamiltonian (9) in the odd bracket

$$\begin{aligned} \dot{x}^i = (x^i, \overset{0}{H})_0 = \{x^i, \overset{1}{H}\}_1 &\equiv x^i \left(\overleftarrow{\partial}_q \overrightarrow{\partial}_\pi - \overleftarrow{\partial}_\pi \overrightarrow{\partial}_q \right. \\ &\quad \left. - \overleftarrow{\partial}_\pi \overrightarrow{\partial}_\theta + \overleftarrow{\partial}_\theta \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_\theta \right) \overset{1}{H} , \end{aligned} \quad (12)$$

while the equations

$$\dot{\theta} = \pi , \quad \dot{\pi} = -\theta V'' , \quad (13)$$

which are Hamiltonian equations for the odd phase coordinates $\theta^i = (\theta, \pi)$ described in the odd bracket (12)

$$\dot{\theta}^i = \{\theta^i, \overset{1}{H}\}_1$$

can be obtained from (10) with the use of the exterior differential and the map λ

$$dx^i \xrightarrow{\lambda} \theta^i .$$

In relations (11) and (12) $\overleftarrow{\partial}$ and $\overrightarrow{\partial}$ are right and left derivatives, respectively.

Note that the even bracket (11) for arbitrary functions A, B of canonical variables rewritten in terms of the proper time

$$\begin{aligned} t[\overset{0}{H}(q, p), q] &= \int \frac{dq'}{\sqrt{2[\overset{0}{H}(q, p) - V(q')]} \Big|_{q'=q}} \\ &\quad + t_0[\overset{0}{H}(q, p)] \end{aligned}$$

(an arbitrary function t_0 corresponds to the choice of the proper time origin) and its canonically conjugate $\overset{0}{H}$ (8) takes the form

$$(A, B)_0 = A \left(\overleftarrow{\partial}_t \overrightarrow{\partial}_{\overset{0}{H}} - \overleftarrow{\partial}_{\overset{0}{H}} \overrightarrow{\partial}_t \right) B \quad (14)$$

and given in the canonical variables $t, \overset{0}{H}, \overset{1}{H}$ and Θ , where

$$dt \xrightarrow{\lambda} \Theta = \frac{\theta}{p} + \overset{1}{H} \partial_{\overset{0}{H}} t(\overset{0}{H}, q) , \quad (15)$$

the odd bracket (12) for any functions A, B is

$$\begin{aligned} \{A, B\}_1 &= A \left(\overleftarrow{\partial}_t \overrightarrow{\partial}_{\overset{1}{H}} - \overleftarrow{\partial}_{\overset{1}{H}} \overrightarrow{\partial}_t + \overleftarrow{\partial}_\Theta \overrightarrow{\partial}_{\overset{0}{H}} \right. \\ &\quad \left. - \overleftarrow{\partial}_{\overset{0}{H}} \overrightarrow{\partial}_\Theta \right) B . \end{aligned} \quad (16)$$

By using the time derivative and Hamiltonian equations (10), (13), we obtain from the transformations (2) of the configuration superspace variables q and θ the supersymmetry transformation rules for the phase superspace coordinates $z^M = (x^i, \theta^i)$

$$\delta q = i\theta\epsilon , \quad \delta p = i\pi\epsilon ,$$

$$\delta\theta = \epsilon p, \quad \delta\pi = -\epsilon V',$$

which can be uniformly represented by means of the odd bracket (12)

$$\delta z^M = \{z^M, \overset{1}{Q}\}_1 \epsilon$$

with the help of an even supercharge

$$\overset{1}{Q} = \overset{0}{H} + i\pi\theta = \overset{0}{H} + i\overset{1}{H}\Theta, \quad (17)$$

where the last expression follows from (9) and (15).

The even supercharge (17) together with the odd Hamiltonian (9) satisfies in the odd bracket (16) $N = 1$ superalgebra relations

$$\{\overset{1}{Q}, \overset{1}{Q}\}_1 = 2i\overset{1}{H}, \quad \{\overset{1}{Q}, \overset{1}{H}\}_1 = 0.$$

Note that an even covariant derivative has the form

$$\overset{1}{D} = \overset{0}{H} - i\pi\theta = \overset{0}{H} - i\overset{1}{H}\Theta \quad (18)$$

and obeys in the odd bracket (16) the following permutation relations:

$$\{\overset{1}{D}, \overset{1}{D}\}_1 = -2i\overset{1}{H}, \quad \{\overset{1}{D}, \overset{1}{H}\}_1 = 0,$$

$$\{\overset{1}{D}, \overset{1}{Q}\}_1 = 0.$$

4. Superfield approach

The Euler-Lagrange equation

$$\frac{d}{dt}(\partial_{\dot{\Phi}}\mathcal{L}) - \partial_{\Phi}\mathcal{L} = 0$$

for the even Lagrangian density superfield

$$\mathcal{L} = \frac{\dot{\Phi}^2}{2} - V(\Phi), \quad (19)$$

which follows from (4), takes the form

$$\ddot{\Phi} + V'(\Phi) = 0. \quad (20)$$

The components of (20) coincide with equations (6) and (7). From (19) we can define a superfield of the momentum density

$$\mathcal{P} = p + i\eta\pi = \partial_{\dot{\Phi}}\mathcal{L} = \dot{\Phi} = \dot{q} + i\eta\dot{\theta}$$

and a superfield for the Hamiltonian density

$$\mathcal{H} = \overset{0}{H} + i\eta\overset{1}{H} = \dot{\Phi}\mathcal{P} - \mathcal{L} = \frac{\mathcal{P}^2}{2} + V(\Phi). \quad (21)$$

It is remarkable enough that on functions A and B , depending on the real bosonic superfield Φ and the momentum density superfield $\mathcal{P}$, we are able to introduce an *even* Poisson bracket of the following form

$$\{A, B\}_0 = A \left(\overleftarrow{\partial}_{\Phi} \overrightarrow{\partial}_{\mathcal{P}} - \overleftarrow{\partial}_{\mathcal{P}} \overrightarrow{\partial}_{\Phi} \right) B.$$

By means of this bracket with the help of the Hamiltonian density (21), we can determine the following superfield Hamiltonian equations

$$\dot{\Phi} = \{\Phi, \mathcal{H}\}_0 = \mathcal{P},$$

$$\dot{\mathcal{P}} = \{\mathcal{P}, \mathcal{H}\}_0 = -V'(\Phi),$$

which components give Hamiltonian equations (10), (13) for the phase variables $z^M = (q, p; \theta, \pi)$.

5. Conclusion

In the simplest case of the one-dimensional supersymmetric oscillator

$$\overset{1}{H}_{\text{osc}} = p\pi + a^2\theta q, \quad \left(V_{\text{osc}} = \frac{a^2 q^2}{2} \right), \quad (22)$$

where a is a constant of the inverse length dimension, the Hamilton equations for the canonical coordinates $z^M = (q, p; \theta, \pi)$

$$\dot{q} = p, \quad \dot{p} = -a^2 q, \quad \dot{\theta} = \pi, \quad \dot{\pi} = -a^2 \theta$$

can be reproduced both with the odd Hamiltonian (22) in the canonical odd bracket (12) and in the following even Dirac brackets $\{\dots, \dots\}_{0k}^*$, ($k = \pm 1$)

$$\begin{aligned} \dot{z}^M &= \{z^M, \overset{1}{H}_{\text{osc}}\}_1 = \{z^M, \overset{0}{H}_k\}_{0k}^* \\ &\equiv z^M [\overleftarrow{\partial}_q \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_q - ik(\overleftarrow{\partial}_\theta \overrightarrow{\partial}_\pi \\ &\quad + \overleftarrow{\partial}_\pi \overrightarrow{\partial}_\theta)] \overset{0}{H}_k \end{aligned} \quad (23)$$

with the aid of the corresponding even Hamiltonians

$$\overset{0}{H}_k = \frac{p^2 + a^2 q^2}{2} - ik\pi\theta, \quad (24)$$

which coincide in the case with the expressions for the even supercharge $\overset{1}{Q}_{\text{osc}}$ (17) and even covariant derivative $\overset{1}{D}_{\text{osc}}$ (18)

$$\overset{0}{H}_{-1} = \overset{1}{Q}_{\text{osc}} , \quad \overset{0}{H}_1 = \overset{1}{D}_{\text{osc}} .$$

The even Dirac brackets (23) and the even Hamiltonians (24) follow from the even Lagrangians

$$\overset{0}{L}_k = \frac{1}{2} [\dot{q}^2 - a^2 q^2 + ik(\psi^\alpha \dot{\psi}^\alpha + 2a\psi^1 \psi^2)] , \quad (25)$$

where $\psi^\alpha(t)$ ($\alpha = 1, 2$) are two real fermionic coordinates. Indeed, the even Lagrangians (25) lead to the momenta

$$p = \partial_{\dot{q}} \overset{0}{L}_k = \dot{q} , \quad \pi_k^\alpha = \partial_{\dot{\psi}^\alpha} \overset{0}{L}_k = -\frac{ik}{2} \psi^\alpha , \quad (26)$$

canonically conjugate to the coordinates q and ψ^α in the even brackets corresponding to $\overset{0}{L}_k$

$$\{q, p\}_{0k} = 1, \quad \{\psi^\alpha, \pi_k^\beta\}_{0k} = -\delta^{\alpha\beta} . \quad (27)$$

The remaining even-bracket relations between the canonical variables have zero right-hand sides. The last relations in (26) define the second-class constraints

$$\varphi_k^\alpha = \pi_k^\alpha + \frac{ik}{2} \psi^\alpha, \quad \{\varphi_k^\alpha, \varphi_k^\beta\}_0 = -ik\delta^{\alpha\beta}, \quad (28)$$

which commute in the even brackets (27) with the variables

$$\chi_k^\alpha = \pi_k^\alpha - \frac{ik}{2} \psi^\alpha, \quad \{\chi_k^\alpha, \chi_k^\beta\}_0 = ik\delta^{\alpha\beta},$$

$$\{\varphi_k^\alpha, \chi_k^\beta\}_0 = 0 ,$$

entering into the definitions for the even Dirac brackets (23)

$$\begin{aligned} \{z^M, \overset{0}{H}_k\}_{0k}^* &= \{z^M, \overset{0}{H}_k\}_{0k} \\ &- ik\{z^M, \varphi_k^\alpha\}_{0k} \{\varphi_k^\alpha, \overset{0}{H}_k\}_{0k} \\ &= z^M (\overleftarrow{\partial}_q \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_q \\ &\quad + ik \overleftarrow{\partial}_{\chi_k^\alpha} \overrightarrow{\partial}_{\chi_k^\alpha}) \overset{0}{H}_k \end{aligned} \quad (29)$$

and the even Hamiltonians (24)

$$\overset{0}{H}_k = \frac{p^2 + a^2 q^2}{2} + ika\chi_k^1 \chi_k^2 . \quad (30)$$

There is no summation over k in the foregoing formulae. The even Hamiltonians (30) follow from the total even Hamiltonians corresponding to the Lagrangians (25) with the use of the second-class constraints $\varphi_k^\alpha = 0$. A correspondence between different expressions (23), (29) for the even Dirac brackets and (24), (30) for the even Hamiltonians $\overset{0}{H}_k$ can be established by putting either

$$a\chi_k^1 = \pm i\pi, \quad \chi_k^2 = \pm i\theta$$

or

$$a\chi_k^2 = \pm i\pi, \quad \chi_k^1 = \mp i\theta .$$

Note that even in the simplest case of the one-dimensional supersymmetric oscillator the even Lagrangians (25) do not follow from any Lagrangian densities, given in terms of one real scalar bosonic $D = 1, N = 1$ superfield $\Phi(1)^4$, as it takes place for the odd Lagrangian

$$\overset{1}{L}_{\text{osc}} = \dot{q}\dot{\theta} - a^2 q\theta ,$$

that leads to the odd Hamiltonian (22), which is dynamically equivalent (see equation (23)) to the even one (24) followed from the even Lagrangian (25).

Thus, we see that in the case of $D = 1, N = 1$ supersymmetry the description with the odd Lagrangian is *more economical*.

6. Acknowledgements

One of the authors (V.A.S.) would like to thank the administration of the Ludwig Maximilian University of München for hospitality during the period when the main part of this work has been done. V.A.S. is also sincerely grateful to J. Lukierski for kind hospitality at the University of Wrocław where the part of the work has been performed and together with A. Frydryszak for useful discussions.

The research of V.A.S. was supported in part by the Ukrainian State Foundation of Fundamental Researches, Grant No 2.5.1/54 and by Grant INTAS No 93-127 (Extension).

⁴In the case $k = 1$ the even Lagrangian $\overset{0}{L}_1$ (25) is reduced from the even Lagrangian density expressed in terms of one scalar bosonic $D = 1, N = 2$ superfield (see, for example, [6]).

REFERENCES

1. D.A. Leites, Docl. Acad. Nauk SSSR 236 (1977) 804.
2. D.V. Volkov, A.I. Pashnev, V.A. Soroka and V.I. Tkach, JETP Lett. 44 (1986) 70; Theor. Math. Phys. 79 (1989) 424.
3. V.A. Soroka, Lett. Math. Phys. 17 (1989) 201.
4. D.V. Volkov, A.V. Tur, V.V. Yanovsky, Phys. Lett. A 203 (1995) 357.
5. A. Frydryszak, J. Phys. A: Math. Gen. 26 (1993) 7227.
6. V.A. Soroka, Phys. Atom. Nucl. 59 (1996) 1270; hep-th/9507030.
7. A.P. Nersessian, JETP Lett. 58 (1993) 64.